


Satellite and river monitored carbon fluxes in North-eastern Iceland: variations in time and space

Marin Ivanov Kardjilov¹ (marin@hi.is), Gudrun Gisladdottir¹ and Sigurdur Reynir Gislason²

¹ Department of Geology and Geography, Faculty of Science, University of Iceland, Sturlugata 7, IS-101 Reykjavik, Iceland

² Institute of Earth Sciences, University of Iceland, Sturlugata 7, IS-101 Reykjavik, Iceland

Regions dominated by exposed reactive Mg-Ca silicate rocks, high rainfall and high relief are the most important long term CO₂ sinks on land. The basaltic terrain of North-eastern Iceland is one of these regions. The objective of this study is to compare the river carbon fluxes stemming from weathering: dissolved inorganic carbon DIC, dissolved organic carbon, DOC and particulate organic carbon, POC with those of the vegetation in the catchments: gross primary production, GPP and net primary production, NPP. The DIC, DOC and POC fluxes were measured in 8 river catchments in North-eastern Iceland for the 6 years period between 1998 and 2003. The terrestrial GPP and NPP for the same catchments are based on MODIS satellites' data. These remotely sensed fluxes are spatially calculated for every year in the period between 2000 and 2003. The DIC and POC fluxes out of the catchments are studied in relation with the spatial distribution of the vegetation, i.e. with the terrestrial GPP and NPP, temperature and elevation. Thus it was possible to define the climate effect on all the fluxes and the correlation between them.



The 8 catchments cover 1/10th of Iceland and they differ in elevation and bedrock age, vegetation and glacial coverage, precipitation and temperature. Holl and Grimsstadir are the catchments with the highest (11.4 and 10.9 g C m⁻² yr⁻¹) chemical DIC weathering rate while Fjardara is the lowest (3.9 g C m⁻² yr⁻¹) mainly due to the old age of the bedrock. The POC flux for the Holl catchment is highest for all of the years of study (1.0 g C m⁻² yr⁻¹) and the reason is probably the extensive area of wetlands in the catchment, whereas Grimsstadir, Fellsa, Grimsa and Lagarfljot produce the lowest POC fluxes (0.3 g C m⁻² yr⁻¹). The most productive year for riverine fluxes in general is 2002 and the least productive are the years 1998 and 2000 with minor fluctuations for some of the catchments.

The satellite derived GPP and NPP fluxes are highest in the catchments Lagarfljot (92.9 and 23.6 g C m⁻² yr⁻¹) and Grimsa (97.7 and 20.0 g C m⁻² yr⁻¹) and the lowest in Grimsstadir (3.5 and 1.7 g C m⁻² yr⁻¹). The year 2001 is the lowest in terrestrial carbon production while 2003 is the highest one.

The preliminary results indicate correlation between DIC flux and GPP for the catchments of Jokulsa a Dal at Bru (78%) and Jokulsa a Fjollum (55%), small correlation for Jokulsa a Dal at Hjarðarhagi (44%) and Holl (38%) and almost no existence of correlation for the other catchments. DIC correlates better with NPP than GPP for the same catchments, except Holl. POC flux relates to GPP in the similar manner as DIC, but correlates very strong with NPP in the catchments of Jokulsa a Dal at Bru (94%), Jokulsa a Fjollum (83%), Jokulsa a Dal at Hjarðarhagi (64%).

Thus it is evident, that there is no a simple general correlation between the terrestrial carbon fluxes in vegetation and riverine carbon fluxes in the sparsely vegetated catchments of North-eastern Iceland. The vegetation fluxes probably need to reach critical size to strongly affect the carbon fluxes stemming from chemical weathering.

This research was supported by the Icelandic Centre for Research, the Research Fund of the University of Iceland, the National Power Company and the Icelandic Ministry for the Environment.